

arXiv:hep-th/9201027v1 14 Jan 1992

COSET CONSTRUCTIONS IN CHERN-SIMONS GAUGE THEORY

US-FT-15/91
December, 1991

J. M. ISIDRO, J. M. F. LABASTIDA AND A. V. RAMALLO

Departamento de Física de Partículas

Universidad de Santiago

E-15706 Santiago de Compostela, Spain

ABSTRACT

Coset constructions in the framework of Chern-Simons topological gauge theories are studied. Two examples are considered: models of the types $\frac{U(1)_p \times U(1)_q}{U(1)_{p+q}} \cong U(1)_{pq(p+q)}$ with p and q coprime integers, and $\frac{SU(2)_m \times SU(2)_1}{SU(2)_{m+1}}$. In the latter case it is shown that the Chern-Simons wave functionals can be identified with the characters of the minimal unitary models, and an explicit representation of the knot (Verlinde) operators acting on the space of $c < 1$ characters is obtained.

1. *Introduction.* Three years ago Witten found a remarkable equivalence between the Hilbert space of three-dimensional (3D) topological Chern-Simons (CS) theory and the space of conformal blocks of the two-dimensional (2D) Wess-Zumino-Witten (WZW) model with the same gauge group and level [1]. This equivalence has been analyzed and made more explicit in subsequent works by several authors [2,3,4,5,6]. An operator formalism was developed in [5,6] where it was shown that the CS wave functionals can be identified with the characters of the corresponding WZW model. One may ask whether it is essential to have a current algebra in the 2D Rational Conformal Field Theory (RCFT) in order to describe its conformal blocks as wave functionals of a 3D CS gauge theory. As conjectured in [2], Witten's equivalence between the two theories could be extended to any RCFT by using GKO constructions [7] and, therefore, the existence of a current algebra would not be required in general. In this paper we verify this conjecture by showing how the operator formalism constructed in [5,6] can be used to describe coset models. Two examples will be considered. First, we analyze an abelian coset model which will help us to find the good quantization variables for CS coset constructions. Secondly we study the non-abelian $\frac{SU(2)_m \times SU(2)_1}{SU(2)_{m+1}}$ models which are well-known to be equivalent to the minimal unitary models with central charge less than unity [7]. We will be able to formulate in this case a quantum-mechanical problem having the $c < 1$ characters as states and the Verlinde operators [8] associated to any toral knot as observables. In the present letter we give a brief report of our results. An extended version containing details and extensions of our calculations will be presented elsewhere.

2. *Abelian Coset Models.* Let us study the quantization of an abelian topological Chern-Simons (CS) theory defined on a three manifold M with action,

$$S = \frac{p}{2\pi} \int_M A \wedge dA + \frac{q}{2\pi} \int_M B \wedge dB, \quad (1)$$

where p and q are two coprime integers and A and B are $U(1)$ one-form connections. We shall restrict ourselves to manifolds M that can be put at least locally as $\Sigma \times R^1$

where Σ is a compact Riemann surface without boundary, and R^1 will be taken as time direction. On Σ we introduce a system of local complex coordinates z and $\bar{z}$. The corresponding components of the gauge fields parallel to Σ will be denoted as A_z, B_z and $A_{\bar{z}}, B_{\bar{z}}$. In order to proceed to a canonical quantization of the theory let us choose the gauge in which the components of A and B along R^1 vanish. In this gauge the holomorphic components of the gauge fields are canonically conjugate to the antiholomorphic ones in such a way that the only non-vanishing commutation relations on the equal-time surface Σ are,

$$[A_z(\sigma), A_{\bar{z}}(\sigma')] = -\frac{\pi}{2p}\delta^2(\sigma - \sigma'), \quad [B_z(\sigma), B_{\bar{z}}(\sigma')] = -\frac{\pi}{2q}\delta^2(\sigma - \sigma'), \quad (2)$$

where σ^i represent local real coordinates on Σ . When the manifold M has no boundary the action (1) is invariant under gauge transformations: $A \rightarrow A + g^{-1}dg$, $B \rightarrow B + h^{-1}dh$, with $g = e^\alpha$ and $h = e^\beta$. In our gauge these transformations are generated by the operator

$$Q(\alpha, \beta) = -\frac{2}{\pi} \int_{\Sigma} (\alpha p F_{z\bar{z}}[A] + \beta q F_{z\bar{z}}[B]), \quad (3)$$

where $F_{z\bar{z}}[A]$ and $F_{z\bar{z}}[B]$ are field strength tensors (*i.e.*, $F_{z\bar{z}}[A] = \partial_z A_{\bar{z}} - \partial_{\bar{z}} A_z$, etc). The operator that implements gauge transformations in the Hilbert space is simply obtained by exponentiating the generator (3): $G(\alpha, \beta) = e^{Q(\alpha, \beta)}$. The physical states are those left invariant by the action of $G(\alpha, \beta)$. This is the Gauss law, which is enough to determine the states of the theory. From the commutation relations (2) we conclude that one can represent these states as wave functionals depending on the antiholomorphic components of the gauge fields: $\Psi[A_{\bar{z}}, B_{\bar{z}}]$. As A and B are independent fields, the solution to the Gauss law, $G(\alpha, \beta)\Psi = \Psi$, can be factorized as,

$$\Psi[A_{\bar{z}}, B_{\bar{z}}] = \Phi_p[A_{\bar{z}}] \Phi_q[B_{\bar{z}}]. \quad (4)$$

Using the commutation relations (2) and the explicit form of the generator (3), the

Gauss law implies the following equation for $\Phi_p[A_{\bar{z}}]$:

$$\Phi_p[A_{\bar{z}} + \partial_{\bar{z}}\alpha] = e^{\left(-\frac{p}{\pi} \int_{\Sigma} \partial_z \alpha \partial_{\bar{z}} \alpha - \frac{2p}{\pi} \int_{\Sigma} \partial_z \alpha A_{\bar{z}}\right)} \Phi_p[A_{\bar{z}}]. \quad (5)$$

($\Phi_q[B_{\bar{z}}]$ satisfies a similar equation). In order to solve (5), let us restrict ourselves to the case in which Σ is a torus T^2 with modular parameter τ . The holomorphic one-form $\omega(z)$ is defined by means of its integrals along the cycles **A** and **B** of the canonical homology basis: $\int_{\mathbf{A}} \omega = 1$, $\int_{\mathbf{B}} \omega = \tau$. Our conventions are such that the **A**-cycle is contractible in the solid torus and $\int_{T^2} d^2\sigma \omega(z) \overline{\omega(z)} = \tau_2$, with $\tau_2 = \text{Im}\tau$. A general gauge transformation $g : T^2 \rightarrow U(1)$ can be parametrized as $g(z) = \exp[\alpha(z) + \alpha_{mn}(z)]$, where $\alpha(z)$ is a single-valued function on T^2 , whereas $e^{\alpha_{mn}}$ is a function that winds m (n) times around the $U(1)$ group as we move along the **A** (**B**) cycle on T^2 . These large gauge transformations can be parametrized in general with the help of the holomorphic one-forms,

$$\alpha_{mn} = -\frac{\pi}{\tau_2}(n + m\bar{\tau}) \int^z \omega(z) + \frac{\pi}{\tau_2}(n + \tau m) \int^{\bar{z}} \overline{\omega(z)}. \quad (6)$$

We are now ready to obtain the general solution to the Gauss law. Let us first parametrize $A_{\bar{z}}$ and A_z as $A_{\bar{z}} = \partial_{\bar{z}}\chi^A + \frac{\pi\bar{\omega}}{\tau_2}a$, $A_z = -A_{\bar{z}}^\dagger$ where χ^A is a single-valued function and a is a constant complex number containing all the information about the holonomy of the gauge field around the non-trivial cycles of T^2 . Under the gauge transformation induced by g , the variables χ^A and a change as follows: $\chi^A \rightarrow \chi^A + \alpha$, $a \rightarrow a + n + \tau m$. The general solution to the Gauss law (5) can be easily expressed in terms of χ^A and a . It is a linear combination of the functions:

$$\Phi_{j,p}[A_{\bar{z}}] = e^{-\frac{p}{\pi} \int_{T^2} \partial_z \chi^A \partial_{\bar{z}} \chi^A} \varphi_{j,p}(a, \tau), \quad (7)$$

where j is an integer and $\varphi_{j,p}(a, \tau)$ is given by

$$\varphi_{j,p}(a, \tau) = [\eta(\tau)]^{-1} e^{\pi a^2 \tau_2^{-1} p} \theta\left[\frac{j}{0}\right]_{\frac{2p}{0}}(2pa|2p\tau). \quad (8)$$

In (8) $\eta(\tau)$ is the Dedekind η -function, and $\theta\left[\frac{a}{b}\right](z|\tau)$ denotes the Jacobi theta function with characteristics [9]. Notice that $\varphi_{j+2p,p} = \varphi_{j,p}$, and therefore there

are only $2p$ independent solutions to the Gauss law equation (5). Obviously, the general form of $\Phi_q[B_{\bar{z}}]$ can be obtained in the same way. If we parametrize $B_{\bar{z}}$ as $B_{\bar{z}} = \partial_{\bar{z}}\chi^B + \frac{\pi\bar{\omega}}{\tau_2}b$, the $2q$ independent solutions $\Phi_{k,q}[B_{\bar{z}}]$ are given by eqs. (7) and (8) with the changes $j \rightarrow k$, $p \rightarrow q$, $\chi^A \rightarrow \chi^B$ and $a \rightarrow b$. From (7) and (8) it is evident that our states can be identified with the characters of a RCFT with an abelian Kac-Moody symmetry. In the case at hand the basis of our Hilbert space are the characters of a $U(1)_p \times U(1)_q$ current algebra. This is nothing but a particular case of Witten's correspondence between the Hilbert space of CS theories and the space of conformal blocks of RCFT's.

In a path integral approach it is possible to give a representation to the solutions of the Gauss law as functional integrals over gauge field configurations. Suppose that our three-dimensional manifold is a solid torus whose boundary is the torus T^2 on which our wave functionals are defined. We can represent $\Psi[A_{\bar{z}}, B_{\bar{z}}]$ as,

$$\Psi[A_{\bar{z}}, B_{\bar{z}}] = \int [DADB] W_{j,p}^{\gamma}(A) W_{k,q}^{\gamma'}(B) \exp \left[iS - \frac{1}{\pi} \int_{T^2} d^2\sigma (pA_z A_{\bar{z}} + qB_z B_{\bar{z}}) \right]. \quad (9)$$

In (9) we integrate over gauge field configurations on the solid torus having fixed values of $A_{\bar{z}}$ and $B_{\bar{z}}$ on the boundary T^2 . We have also inserted in our functional integral Wilson line operators associated to closed curves γ and γ' in the solid torus. $W_{j,p}^{\gamma}(A)$ is given by the equation

$$W_{j,p}^{\gamma}(A) = \exp \left[-j \int_{\gamma} A \right]. \quad (10)$$

Our Gauss law selects flat connections, which means that the path integral (9) only depends on the homotopy class of the curves γ and γ' . If in addition the $U(1)$ charges j and k are integers, then $W_{j,p}^{\gamma}(A)$ and $W_{k,q}^{\gamma'}(B)$ are gauge invariant under large gauge transformations that wind around $U(1)$ as one goes along γ and γ' . Therefore these operators are the natural observables for a topological gauge theory.

The functional integral representation (9) implies the following inner product for wave functionals:

$$(\tilde{\Psi}, \Psi) = \int [DA_z DB_z DA_{\bar{z}} DB_{\bar{z}}] \exp \left[\frac{2}{\pi} \int_{T^2} d^2\sigma (pA_z A_{\bar{z}} + qB_z B_{\bar{z}}) \right] \overline{\tilde{\Psi}(A_{\bar{z}}, B_{\bar{z}})} \Psi(A_{\bar{z}}, B_{\bar{z}}). \quad (11)$$

It can be checked [5,6] that, with the inner product (11), the states $\Psi_{j,k} = \Phi_{j,p} \Phi_{k,q}$ are orthonormal. Moreover, the elements of this basis of our Hilbert space can be obtained by means of very specific insertions in the path integral (9): the state $\Psi_{j,k}$ is created on T^2 by Wilson lines having $U(1)_p$ and $U(1)_q$ charges j and k for curves homotopic to the **B** cycle of T^2 . We get in this way a correspondence between states and observables similar to what happens in RCFT. In order to check this fact let us consider Wilson line operators associated to curves lying completely on T^2 . These curves are the so-called toral knots, and they are characterized by two coprime integers r and s representing the number of times the path of the knot winds around the **A** and **B** cycles respectively. As we noticed above, the effect of inserting one of these operators only depends on the homotopy class in the solid torus of the corresponding path. Of all the possible curves γ and γ' in (9), the toral knots are the most interesting ones in an operator formalism because they can be represented by well-defined operators acting on the Hilbert space of CS wave functionals. Notice first of all that, with the parametrization of the gauge fields A and B on T^2 given above, the Wilson lines for closed curves on the torus only depend on the holonomy parameters a and b (and on their complex conjugates $\bar{a}$ and $\bar{b}$ appearing in the parametrization of A_z and B_z). From the canonical commutation relations (2) (or equivalently from the inner product (11)) we can read off the following commutators:

$$[\bar{a}, a] = \frac{\tau_2}{2\pi p}, \quad [\bar{b}, b] = \frac{\tau_2}{2\pi q}, \quad (12)$$

and therefore $\bar{a}$ and $\bar{b}$ can be represented as,

$$\bar{a} = \frac{\tau_2}{2\pi p} \frac{\partial}{\partial a}, \quad \bar{b} = \frac{\tau_2}{2\pi q} \frac{\partial}{\partial b}. \quad (13)$$

It is now immediate to write down the operator representation of (10) for γ equal to the toral knot (r, s) (*i.e.*, for $\gamma = r\mathbf{A} + s\mathbf{B}$),

$$W_{j,p}^{(r,s)}(A) = \exp \left[\frac{-j\pi}{\tau_2} (r + s\bar{\tau})a + \frac{j}{2p} (r + s\tau) \frac{\partial}{\partial a} \right]. \quad (14)$$

Using well-known properties of the theta functions it is now easily proved [5] that $W_{l,p}^{(0,1)}(A)W_{m,q}^{(0,1)}(B)\Psi_{j,k} = \Psi_{j+l,k+m}$, which implies that Wilson line operators for $\mathbf{B}$ cycles create $U(1)_p \times U(1)_q$ charges. In view of this result it is natural to associate the “vacuum state” $\Psi_{0,0}$ (which has vanishing charges) to the path integral (11) with no Wilson lines inserted. All the independent states of the Hilbert space are obtained by acting on the vacuum with the “creation” operators $W_{l,p}^{(0,1)}(A)$ and $W_{m,q}^{(0,1)}(B)$. These operators are nothing but the Verlinde operators [8], which represent the primary fields of the current algebra on the space of characters. In fact for any r, s in (14) we get a representation of the Verlinde algebra (fusion rules) for the underlying current algebra: $W_{j,p}^{(r,s)}(A)W_{l,p}^{(r,s)}(A) = W_{j+l,p}^{(r,s)}(A)$, and similarly for the B field.

So far we have described the quantization of a topological gauge theory consisting of two independent Chern-Simons fields A and B . We will now show how this system can also be described in terms of new gauge fields that will allow us to perform a coset construction. First of all let us define new one-form connections,

$$C = \frac{pA + qB}{p + q}, \quad D = \frac{A - B}{p + q}, \quad (15)$$

whose inverse relations are:

$$A = C + qD, \quad B = C - pD. \quad (16)$$

We will show below that the quantum mechanical problem of quantizing the $U(1)_p \times U(1)_q$ CS theory given by the action (1) naturally gives rise to the introduction of the gauge connections C and D defined in (15). In fact we shall

prove that the action, observables and states of the theory (1) can be decomposed in terms of a $U(1)_{p+q} \times U(1)_{pq(p+q)}$ CS theory for the gauge fields C and D . First of all notice that, in the gauge in which the time components of the fields A and B vanish, the commutation relations (2), together with (15), imply:

$$[C_z(\sigma), C_{\bar{z}}(\sigma')] = -\frac{\pi}{2(p+q)}\delta^2(\sigma-\sigma'), \quad [D_z(\sigma), D_{\bar{z}}(\sigma')] = -\frac{\pi}{2pq(p+q)}\delta^2(\sigma-\sigma'). \quad (17)$$

Moreover C and D commute and therefore they can be considered as independent fields. Equation (17) seems to indicate that C and D describe $U(1)$ theories with levels $p+q$ and $pq(p+q)$ respectively. This is indeed the case, as can be seen from the form that the action S given in (1) takes in these variables,

$$S = \frac{(p+q)}{2\pi} \int_M C \wedge dC + \frac{pq(p+q)}{2\pi} \int_M D \wedge dD. \quad (18)$$

In fact, the same decomposition also holds for the observables of the theory. Using equation (16) and the canonical commutation relations (17), it is easy to relate Wilson lines in both sets of variables for a given closed curve γ ,

$$W_{j,p}^\gamma(A)W_{k,q}^\gamma(B) = W_{j+k,p+q}^\gamma(C)W_{jq-kp,pq(p+q)}^\gamma(D). \quad (19)$$

Notice that the Wilson lines in the variables C and D resulting from the product of two gauge invariant operators $W_{j,p}^\gamma(A)W_{k,q}^\gamma(B)$ in (19) have integer charges and therefore are also invariant under large gauge transformations of the fields C and D . They are thus good observables for the theory defined by the action (18) (*i.e.*, when C and D are considered as fundamental fields).

The Hilbert space of states of the $U(1)_p \times U(1)_q$ theory can also be decomposed in terms of $U(1)_{p+q} \times U(1)_{pq(p+q)}$ CS wave functionals. The following relation holds:

$$\Phi_{j,p}[A_{\bar{z}}]\Phi_{k,q}[B_{\bar{z}}] = \sum_{l \in \mathbb{Z}_{p+q}} \Phi_{j+k+2pl,p+q}[C_{\bar{z}}]\Phi_{jq-kp+2pql,pq(p+q)}[D_{\bar{z}}]. \quad (20)$$

In (20) $\Phi_{i,p+q}[C_{\bar{z}}]$ and $\Phi_{i,pq(p+q)}[D_{\bar{z}}]$ are given by equation (7) with the appropriate changes in the arguments. If p and q are coprimes all the $2(p+q)$ ($2pq(p+q)$)

states $\Phi_{i,p+q}[C_{\bar{z}}]$ ($\Phi_{i,pq(p+q)}[D_{\bar{z}}]$) appear on the right-hand side of (20) when the different $U(1)_p \times U(1)_q$ states are decomposed. In order to prove (20) it is enough to check it for the case $j = k = 0$. The proof for the other cases follows from the Wilson line decomposition law (19) and from the fact that in CS theory the Wilson line operators for the **B** homology cycle create the corresponding character when acting on the vacuum. For $j = k = 0$ in (20) we need to study the following product:

$$\theta(2pa|2p\tau)\theta(2qb|2q\tau) = \sum_{m \in Z} \sum_{n \in Z} e^{i\pi\tau(2pm^2+2qn^2)} e^{2\pi i(2pma+2qnb)}, \quad (21)$$

where we have denoted by $\theta(z|\tau)$ the theta functions with vanishing characteristics. Suppose we parametrize C and D on T^2 in a way similar to what we did for A and B and let us denote by c and d the holonomy parameters for C and D respectively. The right-hand side of (21) can be rewritten as a sum of theta functions with c and d in their arguments if one introduces the new summation indices r and s ,

$$m - n = (p + q)s, \quad pm + qn = (p + q)r. \quad (22)$$

Notice that in general r and s are not integers. Let us obtain their general form. From (22) follows that $m = r + qs$, $n = r - ps$, and therefore r and s satisfy the constraints $r + qs \in Z$ and $r - ps \in Z$. In order to solve these constraints let us first parametrize r and s in the form dictated by (22),

$$r = \bar{r} + \frac{k}{p+q}, \quad s = \bar{s} + \frac{l}{p+q}. \quad (23)$$

Now $\bar{r}$ and $\bar{s}$ are independent integers and $k, l \in Z_{p+q}$. The constraints on r and s translate into the simple condition $k = pl \pmod{p+q}$, which solves for k in terms of l . The sums in $\bar{r}$ and $\bar{s}$ build theta functions in the variables c and d respectively.

The final result is,

$$\begin{aligned} & \theta(2pa|2p\tau)\theta(2qb|2q\tau) = \\ & \sum_{l \in Z_{p+q}} \theta \left[\begin{matrix} \frac{2pl}{2(p+q)} \\ 0 \end{matrix} \right] (2(p+q)c|2(p+q)\tau) \theta \left[\begin{matrix} \frac{2pql}{2pq(p+q)} \\ 0 \end{matrix} \right] (2pq(p+q)d|2pq(p+q)\tau) , \end{aligned} \quad (24)$$

which is the crucial equation needed to prove (20) for $j = k = 0$.

From eqs. (18), (19) and (20) we conclude that, as indicated above, C and D are the natural variables to decompose a $U(1)_p \times U(1)_q$ CS theory in terms of states with well-defined quantum numbers under $U(1)_{p+q} \times U(1)_{pq(p+q)}$. Plugging this decomposition into our Hilbert space inner product (11) and integrating out the C field, we end up with an effective quantum-mechanical problem for the remaining field D . Using (16) we obtain that the measure for the inner product (11) is $\exp \left[\frac{2}{\pi} \int d^2\sigma ((p+q)C_z C_{\bar{z}} + pq(p+q)D_z D_{\bar{z}}) \right]$. We thus see that the effective problem in D corresponds to an abelian CS theory with level equal to $pq(p+q)$. The integration of the C field is the CS version of the standard coset procedure of modding out the degrees of freedom corresponding to C . In the case at hand we are performing the coset $\frac{U(1)_p \times U(1)_q}{U(1)_{p+q}}$. Eqs. (18)-(20) mean that,

$$\frac{U(1)_p \times U(1)_q}{U(1)_{p+q}} \cong U(1)_{pq(p+q)}, \quad (25)$$

which is a well-known result at the level of two-dimensional current algebras[★][2].

In order to understand the implications of the change of variables discussed above, let us remark that according to eq. (20) every $U(1)_p \times U(1)_q$ state splits into a sum of $p+q$ states of $U(1)_{p+q} \times U(1)_{pq(p+q)}$. Such a proliferation of states in the new variables originates from the fact that both types of wave functionals are the solutions to different Gauss laws. When the gauge quantization problem is

★ If p and q are not coprime, then the level of the $U(1)$ coset algebra is $\frac{pq(p+q)}{(p,q)^2}$ where (p,q) is the greatest common divisor of p and q . In this case the gauge fields C and D must be defined as in (15) with the changes $p \rightarrow \frac{p}{(p,q)}$, $q \rightarrow \frac{q}{(p,q)}$.

formulated in terms of C and D as fundamental fields, one expects that the operators $W_{l,p+q}^\gamma(C)$ and $W_{n,pq(p+q)}^\gamma(D)$ be gauge invariant for any integer charges l and n and any closed curve γ . Suppose that we perform abelian gauge transformations in A and B : $A \rightarrow A + g^{-1}dg$, $B \rightarrow B + h^{-1}dh$. Let us denote by n_g and n_h the winding numbers of g and h along γ :

$$n_g = \frac{1}{2\pi i} \int_{\gamma} g^{-1}dg, \quad n_h = \frac{1}{2\pi i} \int_{\gamma} h^{-1}dh. \quad (26)$$

If we require gauge invariance of $W_{l,p+q}^\gamma(C)$ and $W_{n,pq(p+q)}^\gamma(D)$ for any $l, n \in \mathbb{Z}$, we must restrict ourselves to gauge transformations such that certain combinations of n_g and n_h are integers (see eq. (15)):

$$\frac{pn_g + qn_h}{p+q} = r, \quad \frac{n_g - n_h}{p+q} = s, \quad r, s \in \mathbb{Z}. \quad (27)$$

From eq. (27) one gets $n_g = r + qs$, $n_h = r - ps$, which in particular imply that $n_g - n_h = (p+q)s$. Therefore n_g and n_h must be equal modulo $p+q$,

$$n_g = n_h \pmod{p+q}. \quad (28)$$

Remember that when we take Wilson lines for A and B with integer charges as basic observables of our theory, n_g and n_h can be any two independent integers. Therefore we see that when we perform the quantization of the system taking C and D as fundamental fields, we are actually reducing the global gauge symmetry of the physical states and thus the Gauss law becomes less restrictive. This explains why we get more gauge-invariant states in the new variables.

We may ask how one can modify the path integral (9) in such a way that a state with well-defined $U(1)_{p+q} \times U(1)_{pq(p+q)}$ quantum numbers is obtained. In (9) the path integral is performed over A and B and we wish to obtain a state of those created by acting on the vacuum $\Phi_{0,p+q}(C_{\bar{z}})\Phi_{0,pq(p+q)}(D_{\bar{z}})$ with **B**-cycle Wilson line operators for the gauge fields C and D . This situation is reminiscent

of the orbifold constructions [10] in which one describes a system (for example the Ashkin-Teller model) in terms of fields (a scalar field in this example) which are not the basic variables of the system (the spin fields). In order to produce the correct spectrum of these models one has to introduce a projector in the path integral which is equivalent to sum over all possible twisted sectors of the theory. In our case let us show that something similar happens. We first notice that eq. (20) can be inverted. The products of theta-functions appearing in the right-hand side of (20) can be written as,

$$\begin{aligned} & \varphi_{j+k,p+q}(c, \tau) \varphi_{jq-kp,pq(p+q)}(d, \tau) = \\ & = \frac{1}{p+q} \sum_{l \in Z_{p+q}} e^{-\frac{i\pi l(jq-kp)}{pq(p+q)}} e^{\frac{\pi pa^2}{\tau_2}} \theta \left[\begin{matrix} j \\ 2p \\ l \\ p+q \end{matrix} \right] (2pa|2p\tau) \frac{e^{\frac{\pi qb^2}{\tau_2}}}{\eta(\tau)} \theta \left[\begin{matrix} k \\ 2q \\ -l \\ p+q \end{matrix} \right] (2qb|2q\tau). \end{aligned} \quad (29)$$

An important point in analyzing this equation is the fact that the lower characteristics appearing in (29) can be generated by acting with Wilson line operators for the **A**-cycle. Suppose we consider the case $j = k = 0$ (using (19) for a **B**-cycle the analysis can be trivially extended to any j and k). From the operator representation of the Wilson lines (see eq. (14)) and the decomposition law (19) one gets a relation between both types of vacua,

$$\Phi_{0,p+q}(C_{\bar{z}}) \Phi_{0,pq(p+q)}(D_{\bar{z}}) = \left(\frac{1}{p+q} \sum_{l \in Z_{p+q}} W_{l,pq(p+q)}^{(1,0)}(D) \right) \Phi_{0,p}(A_{\bar{z}}) \Phi_{0,q}(B_{\bar{z}}). \quad (30)$$

Let us now define the operator P as follows:

$$P = \frac{1}{p+q} \sum_{l \in Z_{p+q}} W_{l,pq(p+q)}^{(1,0)}(D). \quad (31)$$

It is easy to check that P is a projector when acting on the vacuum $\Phi_{0,p}(A_{\bar{z}}) \Phi_{0,q}(B_{\bar{z}})$ (*i.e.*, $P^2 = P$ and $P^\dagger = P$). Finally eq. (30) allows us to represent the $U(1)_{p+q} \times U(1)_{pq(p+q)}$ vacuum as a path integral over the fields A

and B ,

$$\Phi_{0,p+q}(C_{\bar{z}})\Phi_{0,pq(p+q)}(D_{\bar{z}}) = \int [DADB] P \exp \left[iS - \frac{1}{\pi} \int_{T^2} d^2\sigma (pA_z A_{\bar{z}} + qB_z B_{\bar{z}}) \right]. \quad (32)$$

The extension of this result for arbitrary charges j and k is straightforward provided the adequate Wilson line operators are introduced in the functional integral.

3. Non-Abelian Coset Models. Let us now extend the previous formalism to a non-abelian case. Our aim is to construct the CS coset $\frac{SU(2)_m \times SU(2)_1}{SU(2)_{m+1}}$ which gives rise [7] at the two-dimensional level to the minimal unitary representations of the Virasoro algebra with central charge $c_m = 1 - \frac{6}{(m+2)(m+3)}$, for $m = 1, 2, \dots$ [10]. Our starting point will be the non-abelian CS theory for two independent gauge fields A and B having levels m and 1 respectively,

$$S = \frac{m}{4\pi} \int_M Tr \left[A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right] + \frac{1}{4\pi} \int_M Tr \left[B \wedge dB + \frac{2}{3} B \wedge B \wedge B \right]. \quad (33)$$

As was proved in [5,6], the operator formalism of a non-abelian CS theory on T^2 gives rise to an effective quantum-mechanical problem in the holonomy part of the gauge connection, which takes values in the Cartan subalgebra of the gauge group. For an $SU(2)$ gauge field A on the torus, after taking the gauge $A_0 = 0$, this part of the connection can be parametrized as $A_{\bar{z}} = \frac{\pi a}{2\tau_2} \bar{\omega} \sigma_3$, and $A_z = -A_{\bar{z}}^\dagger$, where σ_3 is a Pauli matrix and we are in the gauge $A_0 = 0$. The states that solve the effective Gauss law are of the form [5,6],

$$\Phi_{j,m} = \frac{\lambda_{j,m+2}(a)}{\Pi(a)}, \quad (34)$$

being

$$\lambda_{j,m+2}(a) = e^{\frac{\pi(m+2)a^2}{4\tau_2}} \left[\Theta_{j+1,m+2}(a, \tau, 0) - \Theta_{-j-1,m+2}(a, \tau, 0) \right], \quad (35)$$

where $\Theta_{j,k}$ are classical theta-functions [9,11] and $\Pi(a) = \lambda_{0,2}(a)$. The wave functionals $\Phi_{j,m}(a)$ represent the characters of an $SU(2)$ Kac-Moody algebra at level

m for an isospin $j/2$ [11,12]. Using the periodicity properties of the theta-functions one can check that there are $m+1$ independent states labeled by $j = 0, \dots, m$. The wave functionals (34) are symmetric under the Weyl reflection $a \rightarrow -a$, although both the numerator and the denominator in (34) are antisymmetric. The inner product measure in the effective Hilbert space is $\frac{dad\bar{a}}{2\sqrt{\tau_2}}\Pi(a)\bar{\Pi}(a)e^{\frac{-(m+2)}{2\tau_2}a\bar{a}}$. As the Π factors coming from the measure and the states cancel in the inner product, we can ignore them everywhere and take the numerator of (34) as wave function. In this basis of states the operator formalism greatly simplifies [5,6] since the basic commutator is just $[\bar{a}, a] = \frac{2\tau_2}{\pi(m+2)}$. The observables of the theory are Wilson lines associated to representations of $SU(2)$, $W_{j,m}^\gamma(A) = \text{Tr}_j \exp[-\int_\gamma A]$, the trace taken in the representation of isospin $j/2$ of $SU(2)$. Let us denote by Λ_j the set of weights of the $SU(2)$ irreducible representation of isospin $\frac{j}{2}$ (*i.e.*, the eigenvalues of the generator of the Cartan subalgebra). In our conventions $\Lambda_j = \{-j, -j+2, \dots, j-2, j\}$. The $SU(2)$ operators $W_{j,m}^\gamma(A)$ can be written as a sum of abelian Wilson lines whose charges are the set of weights Λ_j . For a toral knot $\gamma = r\mathbf{A} + s\mathbf{B}$ one gets the operator $W_{j,m}^{(r,s)}(A) = \sum_{n \in \Lambda_j} \exp[\frac{-n\pi}{2\tau_2}(r + s\bar{\tau})a + \frac{n}{m+2}(r + s\tau)\frac{\partial}{\partial a}]$.

Consider now the quantization of the B field in (33). The level one case ($m = 1$) is somehow special. It can be easily shown that for $m = 1$ (and $a \rightarrow b$) we can absorb in (34) the Π factors of the denominator and write the character in the form [13,14],

$$\Phi_{l,1}(b) = \frac{e^{\frac{\pi b^2}{4\tau_2}} \theta\left[\begin{smallmatrix} \frac{l}{2} \\ 0 \end{smallmatrix}\right](b|2\tau)}{\eta(\tau)}, \quad l = 0, 1. \quad (36)$$

Therefore, in this case, the $SU(2)_1$ characters can be written as $U(1)_1$ characters. Accordingly, the Verlinde operators that act correctly on the states (36) can be represented by abelian Wilson lines $W_{l,1}^\gamma(B) = \exp[-l \int_\gamma B]$ (notice that there is no trace!) with $B = \frac{\pi b}{2\tau_2}\bar{\omega} - \frac{\pi \bar{b}}{2\tau_2}\omega$ and now we have the basic commutator $[\bar{b}, b] = \frac{2\tau_2}{\pi}$ (which can be obtained by looking at the measure in the space of level-one abelian characters).

Using the above results we can try to follow the spirit of the abelian coset

construction for this $SU(2)$ case. First of all, in complete agreement with (15), let us define new variables c and d as follows:

$$c = \frac{(m+2)a+b}{m+3}, \quad d = \frac{a-b}{m+3}. \quad (37)$$

Notice that using the commutation relations for the a and b variables one gets $[\bar{c}, c] = \frac{2\tau_2}{\pi(m+3)}$, which suggests that c is a good variable for $SU(2)_{m+1}$ wave functionals. We have learned from our abelian example that, in order to extract $SU(2)_{m+1}$ wave functions from $SU(2)_m \times SU(2)_1$ states, one has to introduce a projector. In the non-abelian case the Weyl group plays a major role. In the basis we have chosen to describe our states (the λ 's in (34)) the wave functions are Weyl antisymmetric. As we argued above c is the natural variable for $SU(2)_{m+1}$ characters. It is thus clear that it will only be possible to decompose into $SU(2)_{m+1}$ states that sector of the $SU(2)_m \times SU(2)_1$ Hilbert space which is antisymmetric under $c \rightarrow -c$. Let us call $\mathcal{P}$ the operator that acting on any function $f(c)$ gives $(f(c) - f(-c))/2$. Following steps similar to those that led us to (20) we get,

$$\mathcal{P}(\lambda_{j,m+2}(a)\Phi_{l,1}(b)) = \sum_{i=0}^{m+1} \lambda_{i,m+3}(c)\chi_{j+1,i+1}(d), \quad (38)$$

where the $\lambda_{i,m+3}(c)$ are given by (35) with $m \rightarrow m+1$ and $a \rightarrow c$. The sum is performed over $j-i$ even (odd) for $l=0$ ($l=1$). The functions $\chi_{p,q}(d)$ are

$$\chi_{p,q}(d, \tau) = \frac{e^{\frac{\pi k d^2}{4\tau_2}}}{2\eta(\tau)} \left[\Theta_{n_-,k}(d, \tau, 0) - \Theta_{n_+,k}(d, \tau, 0) + \Theta_{-n_-,k}(d, \tau, 0) - \Theta_{-n_+,k}(d, \tau, 0) \right], \quad (39)$$

where $n_{\pm} = p(m+3) \pm q(m+2)$, and $k = (m+2)(m+3)$. The wave functions (39) evaluated at the origin ($d=0$) are the familiar Rocha-Caridi [15] characters for the minimal unitary models with central charge $c_m = 1 - \frac{6}{(m+2)(m+3)} < 1$. Notice that at $d=0$ the last two terms of (39) equal the first two ones. From (38) we see that the range of p (q) in $\chi_{p,q}$ is $1 \leq p \leq m+1$ ($1 \leq q \leq m+2$). These

values define the so-called conformal grid. On the other hand, different values of p, q in the conformal grid give rise to the same character $\chi_{p,q}$. In fact, if we change $p \rightarrow m+2-p$ and $q \rightarrow m+3-q$, the $n_{\pm}$ change as $n_+ \rightarrow -n_+ + 2k$, $n_- \rightarrow -n_-$. From our expression (39) we get $\chi_{p,q} = \chi_{m+2-p, m+3-q}$. This is the well-known reflection property of the conformal grid [10]. Notice that at $d \neq 0$ it is essential to have the four terms in (39) in order to fulfill this property. Finally, we can check that our wave functions $\chi_{p,q}$ behave under the modular group exactly as the characters of the minimal models,

$$\begin{aligned}\chi_{p,q}(d, \tau)|_T &\equiv \chi_{p,q}(d, \tau + 1) = e^{2\pi i(h_{p,q} - \frac{cm}{24})} \chi_{p,q}(d, \tau), \\ \chi_{p,q}(d, \tau)|_S &\equiv \chi_{p,q}\left(\frac{d}{\tau}, \frac{-1}{\tau}\right) = \sum_{p', q'} S_{p,q}^{p', q'} \chi_{p', q'}(d, \tau),\end{aligned}\tag{40}$$

where $h_{p,q} = \frac{[(m+3)p - (m+2)q]^2 - 1}{4(m+2)(m+3)}$ are the conformal weights of the primary fields and

$$S_{p,q}^{p', q'} = \left(\frac{8}{(m+2)(m+3)}\right)^{\frac{1}{2}} (-1)^{(p+q)(p'+q')} \sin \frac{\pi p p'}{m+2} \sin \frac{\pi q q'}{m+3}.\tag{41}$$

The above results imply that the states of the effective problem in d (*i.e.*, of the $\frac{SU(2)_m \times SU(2)_1}{SU(2)_{m+1}}$ CS coset theory) are in one-to-one correspondence with the characters of the $c_m < 1$ unitary minimal models. We have thus reproduced the GKO construction of these models within the CS context.

In order to complete our effective quantum-mechanical problem for the coset model we have to find the observables that act on the space of characters (39). These observables must be represented by topologically invariant operators. The obvious choice are Wilson line operators associated to homotopy classes of curves on T^2 . First of all we must define a one-form gauge connection for the variable d . In complete analogy with what we have done for the other variables, let us define the abelian one-form $D = \frac{\pi d}{2\tau_2} \bar{\omega} - \frac{\pi \bar{d}}{2\tau_2} \omega$. From the definition (37) and the commutation relations of a and b we get that $[\bar{d}, d] = \frac{2\tau_2}{\pi(m+2)(m+3)}$, which implies the operator representation $\bar{d} = \frac{2\tau_2}{\pi(m+2)(m+3)} \frac{\partial}{\partial d}$. For a given closed curve we will

associate observables constructed by combining abelian Wilson line operators for the gauge field D with different charges. We do not want these operators to take us out of the Hilbert space spanned by the states (39). Therefore we must choose the charges entering into the combination in such a way that the resulting operator carries the quantum numbers of the characters $\chi_{p,q}$. In this way we will get an association between states and operators which is to be expected for a topological gauge theory. A glance at equation (38) reveals the origin of the quantum numbers p and q in $\chi_{p,q}$: $\frac{p-1}{2}$ is the $SU(2)_m$ isospin of the initial state (before any projection) whereas $\frac{q-1}{2}$ is the isospin of the $SU(2)_{m+1}$ state that we mod out when we perform the coset construction. Therefore it is natural to suspect that the $U(1)$ charges needed to build up Virasoro quantum numbers can be obtained by combining the weights of two $SU(2)$ representations of isospins $\frac{p-1}{2}$ and $\frac{q-1}{2}$. Let us check that this is indeed the case. Define the set of weights $\Gamma_{p,q} = (m+3)\Lambda_{p-1} + (m+2)\Lambda_{q-1}$ and the operators $W_{p,q}^\gamma = \sum_{n \in \Gamma_{p,q}} e^{-n \int_\gamma D}$. For a toral knot $\gamma = r\mathbf{A} + s\mathbf{B}$ we can write the explicit representation of $W_{p,q}^{(r,s)}$:

$$W_{p,q}^{(r,s)} = \sum_{n \in \Gamma_{p,q}} \exp\left[-\frac{n\pi}{2\tau_2}(r + s\bar{\tau})d + \frac{n}{(m+2)(m+3)}(r + s\tau)\frac{\partial}{\partial d}\right]. \quad (42)$$

After some calculations we obtain the result of acting with the operator (42) on an arbitrary state,

$$W_{p,q}^{(r,s)} \chi_{p',q'}(d, \tau) = \sum_{n \in \Lambda_{p-1}} \sum_{l \in \Lambda_{q-1}} M_{nl}^{(r,s)}(p, q; p', q') \chi_{p'+sn, q'+sl}(d, \tau), \quad (43)$$

where,

$$M_{nl}^{(r,s)}(p, q; p', q') = e^{i\pi r[s(p-1)(q-1) + (p-1)q' + (q-1)p' + \frac{m+3}{m+2}(\frac{sn^2}{2} + p'n) + \frac{m+2}{m+3}(\frac{sl^2}{2} + q'l)]}. \quad (44)$$

From eq. (44) we confirm that the operators (43) map the Virasoro characters (39) into themselves. In particular from the general expressions given above it is

easy to check that the operators $W_{p,q}^{(0,1)}$ associated to the non-contractible **B**-cycle create the state $\chi_{p,q}$ when acting on the vacuum $\chi_{1,1}$,

$$W_{p,q}^{(0,1)} \chi_{1,1} = \chi_{p,q}. \quad (45)$$

This last equation show us that the operators (42) really carry the quantum numbers associated to the degenerate representations of the Virasoro algebra. As expected the operators associated to the contractible **A**-cycle act diagonally on the space of characters. Their matrix elements are ratios of elements of the modular S matrix,

$$W_{p,q}^{(1,0)} \chi_{p',q'} = \frac{S_{p,q}^{p'q'}}{S_{1,1}^{p'q'}} \chi_{p'q'}. \quad (46)$$

Equations (45) and (46) imply that $W_{p,q}^{(r,s)}$ are the Verlinde operators [8] associated to arbitrary toral knots. Performing modular transformations we can relate operators having different values of r and s in (42). In particular, one can get the **B**-cycle operators by conjugating the operators corresponding to the **A**-cycle with the S matrix (41). This fact, together with the Verlinde theorem [8], ensures that, for a fixed (r, s) toral knot, the operators (43) satisfy the fusion rules of the corresponding primary fields of the 2D conformal field theories.

4. Concluding remarks In this work we have presented a brief account of our main results. There are several possible applications of our approach. For example, from the general matrix element (43) we could compute knot polynomials [16] for the unitary minimal Virasoro models in the same way as they were computed for the WZW model in Ref. [6]. It is also possible in principle to extend our formalism to other coset theories such as, for example, the $N = 1$ supersymmetry minimal models (obtained in [7] from the coset models $\frac{SU(2)_m \times SU(2)_2}{SU(2)_{m+2}}$) and the minimal series of the W_N algebras (described by the coset models $\frac{SU(N)_m \times SU(N)_1}{SU(N)_{m+1}}$). We expect to report on these issues in a future work.

REFERENCES

1. E. Witten, *Comm. Math. Phys.* **121**(1989), 351
2. G. Moore and N. Seiberg, *Phys. Lett.* **B220**(1989), 422
3. S. Elitzur, G. Moore, A. Schwimmer and N. Seiberg, *Nucl. Phys.* **B326**(1989), 108
4. M. Bos and V.P. Nair, *Phys. Lett.* **B223**(1989), 61, *Int. J. Mod. Phys.* **A5**(1990), 959
5. J.M.F. Labastida and A.V. Ramallo, *Phys. Lett.* **B227**(1989), 92, *Phys. Lett.* **228**(1989), 214, *Nucl. Phys. B* (Proc. Suppl.) **16B** (1990) 594
6. J.M.F. Labastida, P.M. Llatas and A.V. Ramallo, *Nucl. Phys.* **B348**(1991), 651
7. P. Goddard, A. Kent and D. Olive, *Phys. Lett.* **B152**(1985), 88, *Comm. Math. Phys.* **103**(1986), 535
8. E. Verlinde, *Nucl. Phys.* **B300**(1988), 360
9. See, for example, D. Mumford, *Tata Lectures on Theta*, Birkhäuser, Basel, 1983
10. See, for example, P. Ginsparg, “Applied Conformal Field Theory”, in *Fields, Strings and Critical Phenomena*, ed. by E. Brezin and J. Zinn-Justin, North Holland, Amsterdam, 1990
11. V.G.Kac, *Infinite Dimensional Lie Algebras*, (Birkhäuser, Basel, 1983)
12. D. Gepner and E. Witten, *Nucl. Phys.* **B278**(1986), 493
13. I.B. Frenkel and V.G. Kac, *Invent. Math.* **62** (1980) 301
14. G. Segal, *Comm. Math. Phys.* **80**(1981), 301
15. A. Rocha-Caridi, in *Vertex Operators in Mathematics and Physics*, ed. by J. Lepowsky, S. Mandelstam and I. Singer, MSRI Publication # 3, Springer, Heidelberg, 1984, 451-473

16. For a review see L. H. Kauffman, *Knots and Physics*, World Scientific, Singapore, 1991, and references therein.